

HCR v3.0 protocol for whole-mount nematode larvae (*Caenorhabditis elegans*)

This protocol has not been optimized for all stages and should only be used as a template.



Technical support: support@molecularinstruments.com

Safety data sheets (SDS): www.molecularinstruments.com/safety

Molecular Instruments products are protected by and for use under patents: www.molecularinstruments.com/patents

Storage conditions

Store HCR probe sets, HCR amplifiers, HCR probe hybridization buffer, and HCR probe wash buffer at -20 °C. Store HCR amplification buffer at 4 °C. On the bench top, keep stock solutions on ice. Make sure all solutions are well mixed before use.



Preparation of fixed whole-mount nematode larvae

1. Wash nematode larvae off plate with 1 mL of M9 buffer and transfer to a 1.5 mL tube.
2. Centrifuge at $200 \times g$ for 2 min to bring larvae to the bottom of the tube and remove the M9 buffer.
3. Wash larvae 3 times with 1 mL of M9 buffer each. Centrifuge at $200 \times g$ for 2 min between washes.
4. Centrifuge and remove $\approx 800 \mu\text{L}$ of M9 buffer.
5. Aliquot larvae sufficient for a reaction into 1.5 mL tubes.
6. Add 1 mL of 4% paraformaldehyde (PFA).
CAUTION: use PFA with extreme care as it is a hazardous material.
NOTE: use fresh PFA to avoid increased autofluorescence.
7. Immediately freeze sample at -80°C overnight before use.
NOTE: Larvae could stay in -80°C freezer for long-term storage.
8. Fix larvae by thawing at room temperature for 45 min.
9. Wash larvae 2 times with 1 mL of PBST each. Bring larvae to the bottom of the tube with centrifugation at $200 \times g$ for 2 min in between washes.
10. Treat larvae with 1 mL of proteinase K ($100 \mu\text{g}/\text{mL}$) for 10 min at 37°C .
NOTE: Proteinase K concentration and treatment time should be reoptimized for each batch of proteinase K, or for samples at a different developmental stage.
11. Wash larvae 2 times with 1 mL of PBST each.
12. Incubate in 1 mL of 2 mg/mL of glycine solution for 15 min on ice.
13. Wash larvae 2 times with 1 mL of PBST each.

Buffer recipes for sample preparation

M9 buffer

22 mM KH_2PO_4
42 mM Na_2HPO_4
20.5 mM NaCl
1 mM MgSO_4

For 1 L of solution

3 g of KH_2PO_4
6 g of Na_2HPO_4
5 g of NaCl
1 mL of 1 M MgSO_4
Fill up to 1 L with ultrapure H_2O
Sterilize by autoclaving
Store buffer at 4 °C before use

4% Paraformaldehyde (PFA)

4% PFA
1× PBS

For 40 mL of solution

10 mL of 16% PFA solution
4 mL of 10× PBS
Fill up to 40 mL with ultrapure H_2O

PBST

1× PBS
0.1% Tween 20

For 50 mL of solution

5 mL of 10× PBS
500 μL of 10% Tween 20
Fill up to 50 mL with ultrapure H_2O

Proteinase K solution

100 $\mu\text{g}/\text{mL}$ proteinase K

For 1 mL of solution

5 μL of 20 mg/mL proteinase K
Fill up to 1 mL with PBST

Glycine solution

2 mg/mL glycine
PBST

For 50 mL of solution

100 mg of glycine
Fill up to 50 mL with PBST

NOTE: avoid using calcium chloride and magnesium chloride in PBS as this leads to increased autofluorescence in the samples.

Multiplexed HCR v3.0 protocol

Detection stage

1. Incubate larvae in 1 mL of 50% PBST / 50% probe hybridization buffer for 5 min at room temperature.
CAUTION: probe hybridization buffer contains formamide, a hazardous material.
2. Centrifuge at $200 \times g$ for 2 min to remove solution.
3. Pre-hybridize larvae in 300 μL of probe hybridization buffer at 37°C for 1 h.
4. Prepare probe solution by adding 2 pmol of each probe mixture (e.g. 2 μL of 1 μM stock) to 200 μL of probe hybridization buffer at 37°C .
NOTE: For dHCR imaging, use higher concentration of probe to improve probe hybridization efficiency.
5. Add the probe solution to reach a final hybridization volume of 500 μL .
6. Incubate larvae overnight (12–16 h) at 37°C .
7. Remove excess probes by washing larvae 4×15 min with 1 mL of probe wash buffer at 37°C .
CAUTION: probe wash buffer contains formamide, a hazardous material.
NOTE: pre-heat probe wash buffer to 37°C before use.
NOTE: bring larvae to the bottom of the tube with centrifugation at $500 \times g$ for 2 min for each wash.
8. Wash larvae 2×5 min with 1 mL of $5 \times$ SSCT at room temperature.

Amplification stage

1. Pre-amplify larvae with 300 μL of amplification buffer for 30 min at room temperature.
NOTE: equilibrate amplification buffer to room temperature before use.
2. Separately prepare 30 pmol of hairpin h1 and 30 pmol of hairpin h2 by snap cooling 10 μL of 3 μM stock (heat at 95°C for 90 seconds and cool to room temperature in a dark drawer for 30 min).
NOTE: HCR hairpins h1 and h2 are provided in hairpin storage buffer ready for snap cooling. h1 and h2 should be snap cooled in separate tubes.
3. Prepare hairpin mixture by adding snap-cooled h1 hairpins and snap-cooled h2 hairpins to 200 μL of amplification buffer at room temperature.
4. Add the hairpin mixture to reach a final amplification volume of 500 μL .
5. Incubate the larvae overnight (12–16 h) in the dark at room temperature.
NOTE: For dHCR imaging, amplify for a shorter period of time to ensure single-molecule dots are diffraction-limited.
6. Remove excess hairpins by washing with 1 mL of $5 \times$ SSCT at room temperature:
 - (a) 2×5 min
 - (b) 2×30 min
 - (c) 1×5 min
NOTE: bring larvae to the bottom of the tube with centrifugation at $500 \times g$ for 2 min for each wash.
7. Samples can be stored at 4°C protected from light before microscopy.

Citation Notes

For citation, please select from the list below as appropriate for your application:

- **HCR v3.0**

Automatic background suppression for dramatically enhanced performance (signal-to-background, qHCR precision, dHCR fidelity) and ease-of-use (no probe set optimization for new targets and organisms) (Choi *et al.*, 2018). Quantitative analysis modes:

qHCR imaging: analog mRNA relative quantitation with subcellular resolution in the anatomical context of thick autofluorescent samples.

qHCR flow cytometry: analog mRNA relative quantitation for high-throughput expression profiling of mammalian and bacterial cells.

dHCR imaging: digital mRNA absolute quantitation with single-molecule resolution in the anatomical context of thick autofluorescent samples.

Protocols for v3.0 in diverse organisms are adapted from the Zoo paper.

- **qHCR imaging**

mRNA relative quantitation with subcellular resolution in the anatomical context of whole-mount vertebrate embryos; read-out/read-in analysis framework (Trivedi *et al.*, 2018).

- **Zoo paper**

Protocols for multiplexed mRNA imaging in diverse sample types (Choi *et al.*, 2016):

bacteria in suspension	FFPE human tissue sections
generic sample in suspension	generic sample on slide
whole-mount chicken embryos	whole-mount fruit fly embryos
whole-mount mouse embryos	whole-mount sea urchin embryos
whole-mount worm larvae	whole-mount zebrafish embryos and larvae

- **dHCR imaging**

Single-molecule mRNA imaging in thick autofluorescent samples (e.g., 0.5 mm adult mouse brain sections) (Shah *et al.*, 2016).

- **qHCR northern blot**

Simultaneous quantification of RNA target size and abundance for up to 5 target RNAs (Schwarzkopf & Pierce, 2016).

- **HCR v2.0**

2nd generation in situ HCR technology (v2.0) using DNA HCR probes and DNA HCR amplifiers: 10× increase in signal, 10× reduction in cost, dramatic increase in reagent durability (Choi *et al.*, 2014).

- **HCR v1.0**

1st generation in situ HCR technology (v1.0) using RNA HCR probes and RNA HCR amplifiers: multiplexed mRNA imaging in whole-mount vertebrate embryos with simultaneous signal amplification for up to 5 target mRNAs (Choi *et al.*, 2010).

- **HCR mechanism**

Isothermal enzyme-free molecular signal amplification (Dirks & Pierce, 2004).

References

- Choi, H. M. T., Chang, J. Y., Trinh, L. A., Padilla, J. E., Fraser, S. E., & Pierce, N. A. (2010). Programmable in situ amplification for multiplexed imaging of mRNA expression. *Nat Biotechnol*, **28**(11), 1208–12.
- Choi, H. M. T., Beck, V. A., & Pierce, N. A. (2014). Next-generation in situ hybridization chain reaction: higher gain, lower cost, greater durability. *ACS Nano*, **8**(5), 4284–4294.
- Choi, H. M. T., Calvert, C. R., Husain, N., Huss, D., Barsi, J. C., Deverman, B. E., Hunter, R. C., Kato, M., Lee, S. M., Abelin, A. C. T., Rosenthal, A. Z., Akbari, O. S., Li, Y., Hay, B. A., Sternberg, P. W., Patterson, P. H., Davidson, E. H., Mazmanian, S. K., Prober, D. A., van de Rijn, M., Leadbetter, J. R., Newman, D. K., Readhead, C., Bronner, M. E., Wold, B., Lansford, R., Sauka-Spengler, T., Fraser, S. E., & Pierce, N. A. (2016). Mapping a multiplexed zoo of mRNA expression. *Development*, **143**, 3632–3637.
- Choi, H. M. T., Schwarzkopf, M., Fornace, M. E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha, A., & Pierce, N. A. (2018). Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development*, **145**, dev165753.
- Dirks, R. M., & Pierce, N. A. (2004). Triggered amplification by hybridization chain reaction. *Proc Natl Acad Sci USA*, **101**(43), 15275–15278.
- Schwarzkopf, M., & Pierce, N. A. (2016). Multiplexed miRNA northern blots via hybridization chain reaction. *Nucleic Acids Res*, **44**(15), e129.
- Shah, S., Lubeck, E., Schwarzkopf, M., He, T.-F., Greenbaum, A., Sohn, C. H., Lignell, A., Choi, H. M. T., Gradi-naru, V., Pierce, N. A., & Cai, L. (2016). Single-molecule RNA detection at depth via hybridization chain reaction and tissue hydrogel embedding and clearing. *Development*, **143**, 2862–2867.
- Trivedi, V., Choi, H. M. T., Fraser, S. E., & Pierce, N. A. (2018). Multidimensional quantitative analysis of mRNA expression within intact vertebrate embryos. *Development*, **145**, dev.156869.