

HCR RNA-FISH protocol for mammalian cells on a chambered slide

This protocol has not been validated for all mammalian cell types and should only be used as a template.

Technical support



support@molecularinstruments.com

Safety data sheets (SDS)

www.molecularinstruments.com/safety

Patents

Molecular Instruments® products are protected by and for use under patents: www.molecularinstruments.com/patents



HCR RNA-FISH

Multiplexed, quantitative, high-resolution RNA imaging

Multiplexed Experiment

- Order one HCR RNA-FISH bundle per target RNA

Example 2-Plex Experiment

- HCR bundle for target mRNA1
 - HCR split-initiator probe set: target mRNA1 for use with amplifier B1
 - HCR amplifier: B1-647
 - HCR RNA-FISH buffers: probe hybridization buffer, probe wash buffer, amplification buffer (for use with all bundles)
- HCR bundle for target mRNA2
 - HCR split-initiator probe set: target mRNA2 for use with amplifier B2
 - HCR amplifier: B2-488

Storage conditions

- Store HCR probe sets, HCR amplifiers, HCR probe hybridization buffer, and HCR probe wash buffer at -20 °C.
- Store HCR amplification buffer at 4 °C.
- On the bench top, keep stock solutions on ice.
- Make sure all solutions are well mixed before use.

Preparation of fixed mammalian cells on a chambered slide

1. Coat bottom of each chamber by applying 300 μL of 0.01% poly-D-lysine prepared in cell culture grade H_2O .
NOTE: A volume of 300 μL is sufficient per chamber on an 8-chamber slide.
2. Incubate for at least 30 min at room temperature.
3. Aspirate the coating solution and wash each chamber twice with molecular biology grade H_2O .
4. Plate desired number of cells in each chamber.
5. Grow cells to desired confluency for 24–48 h.
6. Aspirate growth media and wash each chamber with 300 μL of DPBS.
NOTE: avoid using calcium chloride and magnesium chloride in DPBS as this leads to increased autofluorescence.
7. Add 300 μL of 4% formaldehyde to each chamber.
CAUTION: use formaldehyde with extreme care as it is a hazardous material.
8. Incubate for 10 min at room temperature.
9. Aspirate fixative and wash each chamber $2 \times 300 \mu\text{L}$ of DPBS.
10. Aspirate DPBS and add 300 μL of ice-cold 70% ethanol.
11. Permeabilize cells overnight at -20°C .
12. Cells can be stored at -20°C or 4°C until use.

Multiplexed HCR RNA-FISH protocol

Detection stage

1. Aspirate EtOH and air dry samples at room temperature.
2. Wash samples $2 \times 300 \mu\text{L}$ of $2 \times \text{SSC}$.
3. Pre-hybridize samples in $300 \mu\text{L}$ of probe hybridization buffer for 30 min at 37°C .
CAUTION: probe hybridization buffer contains formamide, a hazardous material.
4. Prepare probe solution by adding 1.2 pmol of each probe set (e.g. $1.2 \mu\text{L}$ of $1 \mu\text{M}$ stock) to $300 \mu\text{L}$ of probe hybridization buffer at 37°C .
NOTE: For dHCR imaging, use higher probe concentration (e.g., 16 nM) to increase probe hybridization yield. If desired, this approach can also be used to increase signal for qHCR imaging.
5. Remove the pre-hybridization solution and add the probe solution.
6. Incubate samples overnight (12–16 h) at 37°C .
7. Remove excess probes by washing 4×5 min with $300 \mu\text{L}$ of probe wash buffer at 37°C .
CAUTION: probe wash buffer contains formamide, a hazardous material.
NOTE: Probe wash buffer should be pre-heated to 37°C before use.
8. Wash samples 2×5 min with $5 \times \text{SSCT}$ at room temperature.

Amplification stage

1. Pre-amplify samples in $300 \mu\text{L}$ of amplification buffer for 30 min at room temperature.
2. Separately prepare 18 pmol of hairpin h1 and 18 pmol of hairpin h2 by snap cooling $6 \mu\text{L}$ of $3 \mu\text{M}$ stock (heat at 95°C for 90 seconds and cool to room temperature in a dark drawer for 30 min).
NOTE: HCR hairpins h1 and h2 are provided in hairpin storage buffer ready for snap cooling. h1 and h2 should be snap cooled in separate tubes.
3. Prepare hairpin solution by adding snap-cooled h1 hairpins and snap-cooled h2 hairpins to $300 \mu\text{L}$ of amplification buffer at room temperature.
4. Remove the pre-amplification solution and add the hairpin solution.
5. Incubate samples overnight (12–16 h) in the dark at room temperature.
NOTE: For single-molecule detection, a 45 min amplification time was used to ensure single-molecule dots are diffraction-limited.
6. Remove excess hairpins by washing 5×5 min with $300 \mu\text{L}$ of $5 \times \text{SSCT}$ at room temperature.
7. Aspirate $5 \times \text{SSCT}$ and add $\approx 100 \mu\text{L}$ of antifade mounting reagent.
8. Samples can be stored at 4°C protected from light prior to imaging.

Buffer recipes

4% formaldehyde in PBS

4% formaldehyde
1× PBS

For 10 mL of solution

2.5 mL of 16% formaldehyde
1 mL of 10× PBS
Fill up to 10 mL with molecular biology grade H₂O

5× SSCT

5× sodium chloride sodium citrate (SSC)
0.1% Tween 20

For 40 mL of solution

10 mL of 20× SSC
400 μL of 10% Tween 20
Fill up to 40 mL with ultrapure H₂O

NOTE: avoid using calcium chloride and magnesium chloride in PBS as this leads to increased autofluorescence in the samples.

HCR Technology Citation Notes

For citation, please select from the list below as appropriate for your application:

- **HCR IHC + HCR RNA-FISH**

HCR IHC + HCR RNA-FISH enables a unified approach to multiplexed, quantitative, high-resolution protein immunohistochemistry (IHC) and RNA fluorescence in situ hybridization (RNA-FISH), with quantitative 1-step enzyme-free HCR signal amplification performed for all protein and RNA targets simultaneously ([Schwarzkopf et al., 2021](#)).

- **HCR IHC**

HCR IHC enables multiplexed, quantitative, high-resolution protein immunohistochemistry (IHC) in highly autofluorescent samples (e.g., FFPE brain tissue sections) ([Schwarzkopf et al., 2021](#)).

- **HCR RNA-FISH (v3.0)**

Third-generation **HCR RNA-FISH (v3.0)** enables multiplexed, quantitative, high-resolution RNA fluorescence in situ hybridization (RNA-FISH) with **automatic background suppression throughout the protocol** for dramatically enhanced performance (signal-to-background, qHCR precision, dHCR fidelity) and ease-of-use (no probe set optimization for new targets and organisms) ([Choi et al., 2018](#)). Quantitative analysis modes:

- **qHCR RNA imaging**: analog mRNA relative quantitation with subcellular resolution in the anatomical context of thick autofluorescent samples.
- **dHCR RNA imaging**: digital mRNA absolute quantitation with single-molecule resolution in the anatomical context of thick autofluorescent samples.
- **qHCR RNA flow cytometry**: analog mRNA relative quantitation for high-throughput expression profiling of mammalian cells and bacteria.

[Protocols for HCR RNA-FISH \(v3.0\)](#) in diverse organisms are adapted from the Zoo paper.

- **qHCR RNA imaging**

qHCR RNA imaging enables mRNA relative quantitation with subcellular resolution in the anatomical context of thick autofluorescent samples (e.g., whole-mount vertebrate embryos). The **read-out/read-in analysis framework** enables bidirectional quantitative discovery in an anatomical context ([Trivedi et al., 2018](#)).

- **Zoo paper**

Protocols for multiplexed mRNA imaging in diverse sample types ([Choi et al., 2016](#)):

- bacteria in suspension
- FFPE human tissue sections
- generic sample in solution
- generic sample on a slide
- mammalian cells on a slide
- mammalian cells in suspension
- whole-mount chicken embryos
- whole-mount fruit fly embryos
- whole-mount mouse embryos
- whole-mount nematode larvae
- whole-mount sea urchin embryos
- whole-mount zebrafish embryos and larvae

- **dHCR imaging**

[dHCR RNA imaging](#) enables RNA absolute quantitation with single-molecule resolution in the anatomical context of thick autofluorescent samples (e.g., 0.5 mm adult mouse brain sections) ([Shah et al., 2016](#)).

- **qHCR northern blots**

[qHCR northern blots](#) enable multiplexed quantification of RNA target size and abundance for up to 5 target RNAs ([Schwarzkopf & Pierce, 2016](#)).

- **HCR RNA-FISH (v2.0)**

Second-generation in situ HCR RNA-FISH technology (v2.0) using DNA HCR probes and DNA HCR probes and DNA HCR amplifiers: 10× increase in signal, 10× reduction in cost, dramatic increase in reagent durability ([Choi et al., 2014](#)).

- **HCR RNA-FISH (v1.0)**

First-generation HCR RNA-FISH technology (v1.0) using RNA HCR probes and RNA HCR amplifiers: multiplexed mRNA imaging in whole-mount vertebrate embryos with simultaneous signal amplification for up to 5 target mRNAs ([Choi et al., 2010](#)).

- **HCR mechanism**

The [hybridization chain reaction \(HCR\) mechanism](#) enables multiplexed, quantitative, 1-step, isothermal, enzyme-free signal amplification in diverse technological settings ([Dirks & Pierce, 2004](#)).

HCR Technology References

- Choi, H. M. T., Chang, J. Y., Trinh, L. A., Padilla, J. E., Fraser, S. E., & Pierce, N. A. (2010). Programmable in situ amplification for multiplexed imaging of mRNA expression. *Nat Biotechnol*, **28**(11), 1208–12.
- Choi, H. M. T., Beck, V. A., & Pierce, N. A. (2014). Next-generation in situ hybridization chain reaction: higher gain, lower cost, greater durability. *ACS Nano*, **8**(5), 4284–4294.
- Choi, H. M. T., Calvert, C. R., Husain, N., Huss, D., Barsi, J. C., Deverman, B. E., Hunter, R. C., Kato, M., Lee, S. M., Abelin, A. C. T., Rosenthal, A. Z., Akbari, O. S., Li, Y., Hay, B. A., Sternberg, P. W., Patterson, P. H., Davidson, E. H., Mazmanian, S. K., Prober, D. A., van de Rijn, M., Leadbetter, J. R., Newman, D. K., Readhead, C., Bronner, M. E., Wold, B., Lansford, R., Sauka-Spengler, T., Fraser, S. E., & Pierce, N. A. (2016). Mapping a multiplexed zoo of mRNA expression. *Development*, **143**, 3632–3637.
- Choi, H. M. T., Schwarzkopf, M., Fornace, M. E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha, A., & Pierce, N. A. (2018). Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development*, **145**, dev165753.
- Dirks, R. M., & Pierce, N. A. (2004). Triggered amplification by hybridization chain reaction. *Proc Natl Acad Sci USA*, **101**(43), 15275–15278.
- Schwarzkopf, M., & Pierce, N. A. (2016). Multiplexed miRNA northern blots via hybridization chain reaction. *Nucleic Acids Res*, **44**(15), e129.
- Schwarzkopf, M., Liu, M.C., Schulte, S.J., Ives, R., Husain, N., Choi, H.M.T., & Pierce, N. A. (2021). Hybridization chain reaction enables a unified approach to multiplexed, quantitative, high-resolution immunohistochemistry and in situ hybridization. *bioRxiv*, 2021.06.02.446311.
- Shah, S., Lubeck, E., Schwarzkopf, M., He, T.-F., Greenbaum, A., Sohn, C. H., Lignell, A., Choi, H. M. T., Gradinaru, V., Pierce, N. A., & Cai, L. (2016). Single-molecule RNA detection at depth via hybridization chain reaction and tissue hydrogel embedding and clearing. *Development*, **143**, 2862–2867.
- Trivedi, V., Choi, H. M. T., Fraser, S. E., & Pierce, N. A. (2018). Multidimensional quantitative analysis of mRNA expression within intact vertebrate embryos. *Development*, **145**, dev.156869.