

HCR RNA-FISH protocol for whole-mount nematode larvae (*Caenorhabditis elegans*)

This protocol has not been optimized for all stages and should only be used as a template.

Technical support



support@molecularinstruments.com

Safety data sheets (SDS)

www.molecularinstruments.com/safety

Patents

Molecular Instruments® products are protected by and for use under patents: www.molecularinstruments.com/patents



HCR RNA-FISH

Multiplexed, quantitative, high-resolution RNA imaging

Multiplexed Experiment

- Order one HCR RNA-FISH bundle per target RNA

Example 2-Plex Experiment

- HCR bundle for target mRNA1
 - HCR split-initiator probe set: target mRNA1 for use with amplifier B1
 - HCR amplifier: B1-647
 - HCR RNA-FISH buffers: probe hybridization buffer, probe wash buffer, amplification buffer (for use with all bundles)
- HCR bundle for target mRNA2
 - HCR split-initiator probe set: target mRNA2 for use with amplifier B2
 - HCR amplifier: B2-488

Storage conditions

- Store HCR probe sets, HCR amplifiers, HCR probe hybridization buffer, and HCR probe wash buffer at -20 °C.
- Store HCR amplification buffer at 4 °C.
- On the bench top, keep stock solutions on ice.
- Make sure all solutions are well mixed before use.

Preparation of fixed whole-mount nematode larvae

1. Wash nematode larvae off plate with 1 mL of M9 buffer and transfer to a 1.5 mL tube.
2. Centrifuge at $200 \times g$ for 2 min to bring larvae to the bottom of the tube and remove the M9 buffer.
3. Wash larvae 3 times with 1 mL of M9 buffer each. Centrifuge at $200 \times g$ for 2 min between washes.
4. Centrifuge and remove $\approx 800 \mu\text{L}$ of M9 buffer.
5. Aliquot larvae sufficient for a reaction into 1.5 mL tubes.
6. Add 1 mL of 4% paraformaldehyde (PFA).
CAUTION: use PFA with extreme care as it is a hazardous material.
NOTE: use fresh PFA to avoid increased autofluorescence.
7. Immediately freeze sample at -80°C overnight before use.
NOTE: Larvae could stay in -80°C freezer for long-term storage.
8. Fix larvae by thawing at room temperature for 45 min.
9. Wash larvae 2 times with 1 mL of PBST each. Bring larvae to the bottom of the tube with centrifugation at $200 \times g$ for 2 min in between washes.
10. Treat larvae with 1 mL of proteinase K ($100 \mu\text{g}/\text{mL}$) for 10 min at 37°C .
NOTE: Proteinase K concentration and treatment time should be reoptimized for each batch of proteinase K, or for samples at a different developmental stage.
11. Wash larvae 2 times with 1 mL of PBST each.
12. Incubate in 1 mL of 2 mg/mL of glycine solution for 15 min on ice.
13. Wash larvae 2 times with 1 mL of PBST each.

Multiplexed HCR RNA-FISH protocol

Detection stage

1. Incubate larvae in 1 mL of 50% PBST / 50% probe hybridization buffer for 5 min at room temperature.
CAUTION: probe hybridization buffer contains formamide, a hazardous material.
2. Centrifuge at $200 \times g$ for 2 min to remove solution.
3. Pre-hybridize larvae in 300 μL of probe hybridization buffer at 37°C for 1 h.
4. Prepare probe solution by adding 2 pmol of each probe set (e.g. 2 μL of 1 μM stock) to 200 μL of probe hybridization buffer at 37°C .
NOTE: For dHCR imaging, use higher probe concentration (e.g., 16 nM) to increase probe hybridization yield. If desired, this approach can also be used to increase signal for qHCR imaging.
5. Add the probe solution to reach a final hybridization volume of 500 μL .
6. Incubate larvae overnight (>12 h) at 37°C .
7. Remove excess probes by washing larvae 4×15 min with 1 mL of probe wash buffer at 37°C .
CAUTION: probe wash buffer contains formamide, a hazardous material.
NOTE: pre-heat probe wash buffer to 37°C before use.
NOTE: bring larvae to the bottom of the tube with centrifugation at $500 \times g$ for 2 min for each wash.
8. Wash larvae 2×5 min with 1 mL of $5\times$ SSCT at room temperature.

Amplification stage

1. Pre-amplify larvae with 300 μL of amplification buffer for 30 min at room temperature.
NOTE: equilibrate amplification buffer to room temperature before use.
2. Separately prepare 30 pmol of hairpin h1 and 30 pmol of hairpin h2 by snap cooling 10 μL of 3 μM stock (heat at 95°C for 90 seconds and cool to room temperature in a dark drawer for 30 min).
NOTE: HCR hairpins h1 and h2 are provided in hairpin storage buffer ready for snap cooling. h1 and h2 should be snap cooled in separate tubes.
3. Prepare hairpin solution by adding snap-cooled h1 hairpins and snap-cooled h2 hairpins to 200 μL of amplification buffer at room temperature.
4. Add the hairpin solution to reach a final amplification volume of 500 μL .
5. Incubate the larvae overnight (>12 h) in the dark at room temperature.
NOTE: For dHCR imaging, amplify for a shorter period of time to ensure single-molecule dots are diffraction-limited.
6. Remove excess hairpins by washing with 1 mL of $5\times$ SSCT at room temperature:
 - (a) 2×5 min
 - (b) 2×30 min
 - (c) 1×5 min
NOTE: bring larvae to the bottom of the tube with centrifugation at $500 \times g$ for 2 min for each wash.
7. Samples can be stored at 4°C protected from light before microscopy.

Buffer recipes

M9 buffer

22 mM KH_2PO_4
42 mM Na_2HPO_4
20.5 mM NaCl
1 mM MgSO_4

For 1 L of solution

3 g of KH_2PO_4
6 g of Na_2HPO_4
5 g of NaCl
1 mL of 1 M MgSO_4
Fill up to 1 L with ultrapure H_2O
Sterilize by autoclaving
Store buffer at 4 °C before use

4% Paraformaldehyde (PFA)

4% PFA
1× PBS

For 40 mL of solution

10 mL of 16% PFA solution
4 mL of 10× PBS
Fill up to 40 mL with ultrapure H_2O

PBST

1× PBS
0.1% Tween 20

For 50 mL of solution

5 mL of 10× PBS
500 μL of 10% Tween 20
Fill up to 50 mL with ultrapure H_2O

Proteinase K solution

100 $\mu\text{g}/\text{mL}$ proteinase K

For 1 mL of solution

5 μL of 20 mg/mL proteinase K
Fill up to 1 mL with PBST

Glycine solution

2 mg/mL glycine
PBST

For 50 mL of solution

100 mg of glycine
Fill up to 50 mL with PBST

5× SSCT

5× sodium chloride sodium citrate (SSC)
0.1% Tween 20

For 40 mL of solution

10 mL of 20× SSC
400 μL of 10% Tween 20
Fill up to 40 mL with ultrapure H_2O

NOTE: avoid using calcium chloride and magnesium chloride in PBS as this leads to increased autofluorescence in the samples.

HCR Technology Citation Notes

For citation, please select from the list below as appropriate for your application:

- **HCR IHC + HCR RNA-FISH**

[HCR IHC + HCR RNA-FISH](#) enables a unified approach to multiplexed, quantitative, high-resolution protein immunohistochemistry (IHC) and RNA fluorescence in situ hybridization (RNA-FISH), with quantitative 1-step enzyme-free HCR signal amplification performed for all protein and RNA targets simultaneously ([Schwarzkopf et al., 2021](#)).

- **HCR IHC**

[HCR IHC](#) enables multiplexed, quantitative, high-resolution protein immunohistochemistry (IHC) in highly autofluorescent samples (e.g., FFPE brain tissue sections) ([Schwarzkopf et al., 2021](#)).

- **HCR RNA-FISH (v3.0)**

Third-generation [HCR RNA-FISH \(v3.0\)](#) enables multiplexed, quantitative, high-resolution RNA fluorescence in situ hybridization (RNA-FISH) with [automatic background suppression throughout the protocol](#) for dramatically enhanced performance (signal-to-background, qHCR precision, dHCR fidelity) and ease-of-use (no probe set optimization for new targets and organisms) ([Choi et al., 2018](#)). Quantitative analysis modes:

- [qHCR RNA imaging](#): analog mRNA relative quantitation with subcellular resolution in the anatomical context of thick autofluorescent samples.
- [dHCR RNA imaging](#): digital mRNA absolute quantitation with single-molecule resolution in the anatomical context of thick autofluorescent samples.
- [qHCR RNA flow cytometry](#): analog mRNA relative quantitation for high-throughput expression profiling of mammalian cells and bacteria.

[Protocols for HCR RNA-FISH \(v3.0\)](#) in diverse organisms are adapted from the Zoo paper.

- **qHCR RNA imaging**

[qHCR RNA imaging](#) enables mRNA relative quantitation with subcellular resolution in the anatomical context of thick autofluorescent samples (e.g., whole-mount vertebrate embryos). The [read-out/read-in analysis framework](#) enables bidirectional quantitative discovery in an anatomical context ([Trivedi et al., 2018](#)).

- **Zoo paper**

Protocols for multiplexed mRNA imaging in diverse sample types ([Choi et al., 2016](#)):

- bacteria in suspension
- FFPE human tissue sections
- generic sample in solution
- generic sample on a slide
- mammalian cells on a slide
- mammalian cells in suspension
- whole-mount chicken embryos
- whole-mount fruit fly embryos
- whole-mount mouse embryos
- whole-mount nematode larvae
- whole-mount sea urchin embryos
- whole-mount zebrafish embryos and larvae

- **dHCR imaging**

[dHCR RNA imaging](#) enables RNA absolute quantitation with single-molecule resolution in the anatomical context of thick autofluorescent samples (e.g., 0.5 mm adult mouse brain sections) ([Shah et al., 2016](#)).

- **qHCR northern blots**

[qHCR northern blots](#) enable multiplexed quantification of RNA target size and abundance for up to 5 target RNAs ([Schwarzkopf & Pierce, 2016](#)).

- **HCR RNA-FISH (v2.0)**

Second-generation in situ HCR RNA-FISH technology (v2.0) using DNA HCR probes and DNA HCR probes and DNA HCR amplifiers: 10× increase in signal, 10× reduction in cost, dramatic increase in reagent durability ([Choi et al., 2014](#)).

- **HCR RNA-FISH (v1.0)**

First-generation HCR RNA-FISH technology (v1.0) using RNA HCR probes and RNA HCR amplifiers: multiplexed mRNA imaging in whole-mount vertebrate embryos with simultaneous signal amplification for up to 5 target mRNAs ([Choi et al., 2010](#)).

- **HCR mechanism**

The [hybridization chain reaction \(HCR\) mechanism](#) enables multiplexed, quantitative, 1-step, isothermal, enzyme-free signal amplification in diverse technological settings ([Dirks & Pierce, 2004](#)).

HCR Technology References

- Choi, H. M. T., Chang, J. Y., Trinh, L. A., Padilla, J. E., Fraser, S. E., & Pierce, N. A. (2010). Programmable in situ amplification for multiplexed imaging of mRNA expression. *Nat Biotechnol*, **28**(11), 1208–12.
- Choi, H. M. T., Beck, V. A., & Pierce, N. A. (2014). Next-generation in situ hybridization chain reaction: higher gain, lower cost, greater durability. *ACS Nano*, **8**(5), 4284–4294.
- Choi, H. M. T., Calvert, C. R., Husain, N., Huss, D., Barsi, J. C., Deverman, B. E., Hunter, R. C., Kato, M., Lee, S. M., Abelin, A. C. T., Rosenthal, A. Z., Akbari, O. S., Li, Y., Hay, B. A., Sternberg, P. W., Patterson, P. H., Davidson, E. H., Mazmanian, S. K., Prober, D. A., van de Rijn, M., Leadbetter, J. R., Newman, D. K., Readhead, C., Bronner, M. E., Wold, B., Lansford, R., Sauka-Spengler, T., Fraser, S. E., & Pierce, N. A. (2016). Mapping a multiplexed zoo of mRNA expression. *Development*, **143**, 3632–3637.
- Choi, H. M. T., Schwarzkopf, M., Fornace, M. E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha, A., & Pierce, N. A. (2018). Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development*, **145**, dev165753.
- Dirks, R. M., & Pierce, N. A. (2004). Triggered amplification by hybridization chain reaction. *Proc Natl Acad Sci USA*, **101**(43), 15275–15278.
- Schwarzkopf, M., & Pierce, N. A. (2016). Multiplexed miRNA northern blots via hybridization chain reaction. *Nucleic Acids Res*, **44**(15), e129.
- Schwarzkopf, M., Liu, M.C., Schulte, S.J., Ives, R., Husain, N., Choi, H.M.T., & Pierce, N. A. (2021). Hybridization chain reaction enables a unified approach to multiplexed, quantitative, high-resolution immunohistochemistry and in situ hybridization. *Development*, **148**(22), dev199847.
- Shah, S., Lubeck, E., Schwarzkopf, M., He, T.-F., Greenbaum, A., Sohn, C. H., Lignell, A., Choi, H. M. T., Gradinaru, V., Pierce, N. A., & Cai, L. (2016). Single-molecule RNA detection at depth via hybridization chain reaction and tissue hydrogel embedding and clearing. *Development*, **143**, 2862–2867.
- Trivedi, V., Choi, H. M. T., Fraser, S. E., & Pierce, N. A. (2018). Multidimensional quantitative analysis of mRNA expression within intact vertebrate embryos. *Development*, **145**, dev.156869.